

endoret®

technology


EFFICACY IN FACIAL REJUVENATION

EFFICACY AND SAFETY OF PLASMA RICH IN GROWTH FACTORS (PRGF) IN FACIAL SKIN REGENERATION: A RANDOMIZED, CONTROLLED CLINICAL TRIAL WITH HYALURONIC ACID


ENDORET® TECHNOLOGY IMPROVES SKIN HYDRATION AND PATIENT SATISFACTION COMPARED TO HYALURONIC ACID TREATMENTS IN FACIAL REJUVENATION

ENHANCED SKIN HYDRATION

SKIN HYDRATION IMPROVEMENT WITH ENDORET® TECHNOLOGY IS UP TO 1.5 TIMES MORE AFTER 3 MONTHS AND UP TO 4 TIMES MORE AFTER 6 MONTHS OVER HYALURONIC ACID.



IMPROVED PATIENT SATISFACTION



94% OF THE PATIENTS REPORT TO BE SATISFIED OR VERY SATISFIED WITH TECHNOLOGY ENDORET® TREATMENT FOR SKIN HYDRATION

The number of very satisfied patients is 73 % higher in the group treated with ENDORET® TECHNOLOGY

ABSTRACT

Eduardo Anitua, Mikel Sánchez, Rosa Sarabia, Julian Sanz, José Javier Aguirre, Gorka Orive.

PRGF efficacy in facial rejuvenation.

PURPOSE

The present study is the first clinical randomized controlled assay with hyaluronic acid (HA) registered in Europe to show efficacy and safety of PRGF for the treatment of skin ageing.

METHODS

A total of 100 patients of different phototypes (I-IV) and ages (31 -65 y.o.) were recruited for the assay. Randomized patients were treated intradermically with either HA or PRGF. HA was used once every month while PRGF was applied once every two weeks for a total of 3 sessions of each treatment. Two follow up visits were scheduled for each patient group at 3 and 6 months post-treatment. Patient satisfaction was assessed by means of "Liquert scale". Corneometry and patient satisfaction were two of the variables measured at each treatment session and on the 3 and 6 months follow up visits.

RESULTS

Corneometry measurements showed a significant improvement in skin hydration at both 3 and 6 months. Furthermore, the effect of PRGF at 6 months remained 11% higher than initial measurement before treatment while only 3 % improvement was observed after HA treatment. The number of very satisfied patients was 73% higher in the group treated with PRGF.

CONCLUSION

Patients treated with PRGF showed higher skin hydration and referred higher levels of satisfaction than those treated with HA. These effects were already observed during the first month of treatment.